



**MERCER COUNTY
CELINA CITY**
Health Department



PREVENT • PROMOTE • PROTECT

Immunizations for the 2016-2017 School Year
Preschool and Kindergarten

Dear Parents/Guardians:

Kindergarteners are required to have the following vaccines:

The 5th Diphtheria, Tetanus and Pertussis, the 4th polio, the 2nd Measles Mumps Rubella, and the 2nd chickenpox vaccines. Children can receive these vaccines any time after their fourth birthday.

Preschoolers are required to be up to date on immunizations. An up to date preschooler has the following vaccines: 4 Diphtheria, Tetanus, and Pertussis, 3 polio, 1 Measles, Mumps Rubella, 1 chickenpox, 3 Hepatitis B, 2 Hepatitis A, 3-4 Haemophilus vaccines and a yearly influenza vaccine. Each individual child's immunization history should be reviewed.

Vaccines can be administered at some physician offices or at the Health Department.

If you wish to have your vaccines at the Mercer County Health Department, please call 419-586-3251 ext 1270 for an appointment or contact us at healthdept@mccchd.org.

For more information on vaccines go to www.immunize.org or www.cdc.gov

Immunization Appointments are available through the week.

Extra clinic Appointments will be available on the following dates

Tuesday	March 15th	9:30am to 5:30pm
Tuesday	April 19th	9:30am to 5:30pm
Tuesday	May 10th	9:30am to 5:30pm
Tuesday	June 14th	9:30am to 5:30pm

Call 419-586-3251 ext 1270 for an appointment

We are contracted with many insurance carriers and will bill the insurance as a courtesy. Please have your insurance cards available. Please Log into your insurance company's website to determine your individual coverage. Federal or State provided vaccines will not be denied to children 18 and under due to inability to pay.

VACCINE INFORMATION STATEMENT

DTaP Vaccine

What You Need to Know

(Diphtheria,
Tetanus and
Pertussis)

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Diphtheria, tetanus, and pertussis are serious diseases caused by bacteria. Diphtheria and pertussis are spread from person to person. Tetanus enters the body through cuts or wounds.

DIPHTHERIA causes a thick covering in the back of the throat.

- It can lead to breathing problems, paralysis, heart failure, and even death.

TETANUS (Lockjaw) causes painful tightening of the muscles, usually all over the body.

- It can lead to “locking” of the jaw so the victim cannot open his mouth or swallow. Tetanus leads to death in up to 2 out of 10 cases.

PERTUSSIS (Whooping Cough) causes coughing spells so bad that it is hard for infants to eat, drink, or breathe. These spells can last for weeks.

- It can lead to pneumonia, seizures (jerking and staring spells), brain damage, and death.

Diphtheria, tetanus, and pertussis vaccine (DTaP) can help prevent these diseases. Most children who are vaccinated with DTaP will be protected throughout childhood. Many more children would get these diseases if we stopped vaccinating.

DTaP is a safer version of an older vaccine called DTP. DTP is no longer used in the United States.

2 Who should get DTaP vaccine and when?

Children should get 5 doses of DTaP vaccine, one dose at each of the following ages:

- 2 months
- 4 months
- 6 months
- 15–18 months
- 4–6 years

DTaP may be given at the same time as other vaccines.

3 Some children should not get DTaP vaccine or should wait

- Children with minor illnesses, such as a cold, may be vaccinated. But children who are moderately or severely ill should usually wait until they recover before getting DTaP vaccine.
- Any child who had a life-threatening allergic reaction after a dose of DTaP should not get another dose.
- Any child who suffered a brain or nervous system disease within 7 days after a dose of DTaP should not get another dose.
- Talk with your doctor if your child:
 - had a seizure or collapsed after a dose of DTaP,
 - cried non-stop for 3 hours or more after a dose of DTaP,
 - had a fever over 105°F after a dose of DTaP.

Ask your doctor for more information. Some of these children should not get another dose of pertussis vaccine, but may get a vaccine without pertussis, called **DT**.

4 Older children and adults

DTaP is not licensed for adolescents, adults, or children 7 years of age and older.

But older people still need protection. A vaccine called **Tdap** is similar to DTaP. A single dose of Tdap is recommended for people 11 through 64 years of age. Another vaccine, called **Td**, protects against tetanus and diphtheria, but not pertussis. It is recommended every 10 years. There are separate Vaccine Information Statements for these vaccines.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

5**What are the risks from DTaP vaccine?**

Getting diphtheria, tetanus, or pertussis disease is much riskier than getting DTaP vaccine.

However, a vaccine, like any medicine, is capable of causing serious problems, such as severe allergic reactions. The risk of DTaP vaccine causing serious harm, or death, is extremely small.

Mild problems (common)

- Fever (up to about 1 child in 4)
- Redness or swelling where the shot was given (up to about 1 child in 4)
- Soreness or tenderness where the shot was given (up to about 1 child in 4)

These problems occur more often after the 4th and 5th doses of the DTaP series than after earlier doses. Sometimes the 4th or 5th dose of DTaP vaccine is followed by swelling of the entire arm or leg in which the shot was given, lasting 1–7 days (up to about 1 child in 30).

Other mild problems include:

- Fussiness (up to about 1 child in 3)
- Tiredness or poor appetite (up to about 1 child in 10)
- Vomiting (up to about 1 child in 50)

These problems generally occur 1–3 days after the shot.

Moderate problems (uncommon)

- Seizure (jerking or staring) (about 1 child out of 14,000)
- Non-stop crying, for 3 hours or more (up to about 1 child out of 1,000)
- High fever, over 105°F (about 1 child out of 16,000)

Severe problems (very rare)

- Serious allergic reaction (less than 1 out of a million doses)
- Several other severe problems have been reported after DTaP vaccine. These include:
 - Long-term seizures, coma, or lowered consciousness
 - Permanent brain damage.

These are so rare it is hard to tell if they are caused by the vaccine.

Controlling fever is especially important for children who have had seizures, for any reason. It is also important if another family member has had seizures. You can reduce fever and pain by giving your child an *aspirin-free* pain reliever when the shot is given, and for the next 24 hours, following the package instructions.

6**What if there is a serious reaction?****What should I look for?**

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS is only for reporting reactions. They do not give medical advice.

7**The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

8**How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

**Vaccine Information Statement
DTaP Vaccine**

5/17/2007

42 U.S.C. § 300aa-26

Office Use Only



Polio Vaccine

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 What is polio?

Polio is a disease caused by a virus. It enters the body through the mouth. Usually it does not cause serious illness. But sometimes it causes paralysis (can't move arm or leg), and it can cause meningitis (irritation of the lining of the brain). It can kill people who get it, usually by paralyzing the muscles that help them breathe.

Polio used to be very common in the United States. It paralyzed and killed thousands of people a year before we had a vaccine.

2 Why get vaccinated?

Inactivated Polio Vaccine (IPV) can prevent polio.

History: A 1916 polio epidemic in the United States killed 6,000 people and paralyzed 27,000 more. In the early 1950's there were more than 25,000 cases of polio reported each year. Polio vaccination was begun in 1955. By 1960 the number of reported cases had dropped to about 3,000, and by 1979 there were only about 10. The success of polio vaccination in the U.S. and other countries has sparked a world-wide effort to eliminate polio.

Today: Polio has been eliminated from the United States. But the disease is still common in some parts of the world. It would only take one person infected with polio virus coming from another country to bring the disease back here if we were not protected by vaccine. If the effort to eliminate the disease from the world is successful, some day we won't need polio vaccine. Until then, we need to keep getting our children vaccinated.

3 Who should get polio vaccine and when?

IPV is a shot, given in the leg or arm, depending on age. It may be given at the same time as other vaccines.

Children

Children get 4 doses of IPV, at these ages:

- A dose at 2 months
- A dose at 4 months
- A dose at 6-18 months
- A booster dose at 4-6 years

Some "combination" vaccines (several different vaccines in the same shot) contain IPV.

Children getting these vaccines may get one more (5th) dose of polio vaccine. This is not a problem.

Adults

Most adults 18 and older do not need polio vaccine because they were vaccinated as children. But some adults are at higher risk and should consider polio vaccination:

- people traveling to areas of the world where polio is common,
- laboratory workers who might handle polio virus, and
- health care workers treating patients who could have polio.

Adults in these three groups:

- who have **never been vaccinated against polio** should get 3 doses of IPV:
 - Two doses separated by 1 to 2 months, and
 - A third dose 6 to 12 months after the second.
- who have had **1 or 2 doses** of polio vaccine in the past should get the remaining 1 or 2 doses. It doesn't matter how long it has been since the earlier dose(s).
- who have had **3 or more doses** of polio vaccine in the past may get a booster dose of IPV.

Your doctor can give you more information.



4**Some people should not get IPV or should wait.****These people should not get IPV:**

- Anyone with a life-threatening allergy to any component of IPV, including the antibiotics neomycin, streptomycin or polymyxin B, should not get polio vaccine. Tell your doctor if you have any severe allergies.
- Anyone who had a severe allergic reaction to a previous polio shot should not get another one.

These people should wait:

- Anyone who is moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting polio vaccine. People with minor illnesses, such as a cold, may be vaccinated.

Ask your doctor for more information.

5**What are the risks from IPV?**

Some people who get IPV get a sore spot where the shot was given. IPV has not been known to cause serious problems, and most people don't have any problems at all with it.

However, any medicine could cause a serious side effect, such as a severe allergic reaction or even death. The risk of polio vaccine causing serious harm is extremely small.

6**What if there is a serious reaction?****What should I look for?**

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS is only for reporting reactions. They do not give medical advice.

7**The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

8**How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

**Vaccine Information Statement (Interim)
Polio Vaccine**

11/8/2011

42 U.S.C. § 300aa-26

Office Use Only



VACCINE INFORMATION STATEMENT

MMR Vaccine

What You Need to Know

(Measles, Mumps
and Rubella)

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información Sobre Vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Measles, mumps, and rubella are serious diseases. Before vaccines they were very common, especially among children.

Measles

- Measles virus causes rash, cough, runny nose, eye irritation, and fever.
- It can lead to ear infection, pneumonia, seizures (jerking and staring), brain damage, and death.

Mumps

- Mumps virus causes fever, headache, muscle pain, loss of appetite, and swollen glands.
- It can lead to deafness, meningitis (infection of the brain and spinal cord covering), painful swelling of the testicles or ovaries, and rarely sterility.

Rubella (German Measles)

- Rubella virus causes rash, arthritis (mostly in women), and mild fever.
- If a woman gets rubella while she is pregnant, she could have a miscarriage or her baby could be born with serious birth defects.

These diseases spread from person to person through the air. You can easily catch them by being around someone who is already infected.

Measles, mumps, and rubella (MMR) vaccine can protect children (and adults) from all three of these diseases.

Thanks to successful vaccination programs these diseases are much less common in the U.S. than they used to be. But if we stopped vaccinating they would return.

2 Who should get MMR vaccine and when?

Children should get 2 doses of MMR vaccine:

- **First Dose:** 12–15 months of age
- **Second Dose:** 4–6 years of age (may be given earlier, if at least 28 days after the 1st dose)

Some infants younger than 12 months should get a dose of MMR if they are traveling out of the country. (This dose will not count toward their routine series.)

Some adults should also get MMR vaccine: Generally, anyone 18 years of age or older who was born after 1956 should get at least one dose of MMR vaccine, unless they can show that they have either been vaccinated or had all three diseases.

MMR vaccine may be given at the same time as other vaccines.

Children between 1 and 12 years of age can get a “combination” vaccine called MMRV, which contains both MMR and varicella (chickenpox) vaccines. There is a separate Vaccine Information Statement for MMRV.

3 Some people should not get MMR vaccine or should wait.

- Anyone who has ever had a life-threatening allergic reaction to the antibiotic neomycin, or any other component of MMR vaccine, should not get the vaccine. Tell your doctor if you have any severe allergies.
- Anyone who had a life-threatening allergic reaction to a previous dose of MMR or MMRV vaccine should not get another dose.
- Some people who are sick at the time the shot is scheduled may be advised to wait until they recover before getting MMR vaccine.
- Pregnant women should not get MMR vaccine. Pregnant women who need the vaccine should wait until after giving birth. Women should avoid getting pregnant for 4 weeks after vaccination with MMR vaccine.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

- Tell your doctor if the person getting the vaccine:
 - Has HIV/AIDS, or another disease that affects the immune system
 - Is being treated with drugs that affect the immune system, such as steroids
 - Has any kind of cancer
 - Is being treated for cancer with radiation or drugs
 - Has ever had a low platelet count (a blood disorder)
 - Has gotten another vaccine within the past 4 weeks
 - Has recently had a transfusion or received other blood products

Any of these might be a reason to not get the vaccine, or delay vaccination until later.

4 What are the risks from MMR vaccine?

A vaccine, like any medicine, is capable of causing serious problems, such as severe allergic reactions.

The risk of MMR vaccine causing serious harm, or death, is extremely small.

Getting MMR vaccine is much safer than getting measles, mumps or rubella.

Most people who get MMR vaccine do not have any serious problems with it.

Mild problems

- Fever (up to 1 person out of 6)
- Mild rash (about 1 person out of 20)
- Swelling of glands in the cheeks or neck (about 1 person out of 75)

If these problems occur, it is usually within 6-14 days after the shot. They occur less often after the second dose.

Moderate problems

- Seizure (jerking or staring) caused by fever (about 1 out of 3,000 doses)
- Temporary pain and stiffness in the joints, mostly in teenage or adult women (up to 1 out of 4)
- Temporary low platelet count, which can cause a bleeding disorder (about 1 out of 30,000 doses)

Severe problems (very rare)

- Serious allergic reaction (less than 1 out of a million doses)
- Several other severe problems have been reported after a child gets MMR vaccine, including:
 - Deafness
 - Long-term seizures, coma, or lowered consciousness
 - Permanent brain damage

These are so rare that it is hard to tell whether they are caused by the vaccine.

5 What if there is a serious reaction?

What should I look for?

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS is only for reporting reactions. They do not give medical advice.

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

7 How can I learn more?

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement (Interim) MMR Vaccine

4/20/2012

42 U.S.C. § 300aa-26

Office Use Only



Chickenpox Vaccine

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

Chickenpox (also called varicella) is a common childhood disease. It is usually mild, but it can be serious, especially in young infants and adults.

- It causes a rash, itching, fever, and tiredness.
- It can lead to severe skin infection, scars, pneumonia, brain damage, or death.
- The chickenpox virus can be spread from person to person through the air, or by contact with fluid from chickenpox blisters.
- A person who has had chickenpox can get a painful rash called shingles years later.
- Before the vaccine, about 11,000 people were hospitalized for chickenpox each year in the United States.
- Before the vaccine, about 100 people died each year as a result of chickenpox in the United States.

Chickenpox vaccine can prevent chickenpox.

Most people who get chickenpox vaccine will not get chickenpox. But if someone who has been vaccinated does get chickenpox, it is usually very mild. They will have fewer blisters, are less likely to have a fever, and will recover faster.

2 Who should get chickenpox vaccine and when?

Routine

Children who have never had chickenpox should get 2 doses of chickenpox vaccine at these ages:

- | | |
|-----------|--|
| 1st Dose: | 12–15 months of age |
| 2nd Dose: | 4–6 years of age (may be given earlier, if at least 3 months after the 1st dose) |

People 13 years of age and older (who have never had chickenpox or received chickenpox vaccine) should get two doses at least 28 days apart.

Catch-up

Anyone who is not fully vaccinated, and never had chickenpox, should receive one or two doses of chickenpox vaccine. The timing of these doses depends on the person's age. Ask your doctor.

Chickenpox vaccine may be given at the same time as other vaccines.

Note: A "combination" vaccine called **MMRV**, which contains both chickenpox and MMR vaccines, may be given instead of the two individual vaccines to people 12 years of age and younger.

3 Some people should not get chickenpox vaccine or should wait.

- People should not get chickenpox vaccine if they have ever had a life-threatening allergic reaction to a previous dose of chickenpox vaccine or to gelatin or the antibiotic neomycin.
- People who are moderately or severely ill at the time the shot is scheduled should usually wait until they recover before getting chickenpox vaccine.
- Pregnant women should wait to get chickenpox vaccine until after they have given birth. Women should not get pregnant for 1 month after getting chickenpox vaccine.
- Some people should check with their doctor about whether they should get chickenpox vaccine, including anyone who:
 - Has HIV/AIDS or another disease that affects the immune system
 - Is being treated with drugs that affect the immune system, such as steroids, for 2 weeks or longer
 - Has any kind of cancer
 - Is getting cancer treatment with radiation or drugs
- People who recently had a transfusion or were given other blood products should ask their doctor when they may get chickenpox vaccine.

Ask your doctor for more information.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

4**What are the risks from chickenpox vaccine?**

A vaccine, like any medicine, is capable of causing serious problems, such as severe allergic reactions. The risk of chickenpox vaccine causing serious harm, or death, is extremely small.

Getting chickenpox vaccine is much safer than getting chickenpox disease. Most people who get chickenpox vaccine do not have any problems with it. Reactions are usually more likely after the first dose than after the second.

Mild problems

- Soreness or swelling where the shot was given (about 1 out of 5 children and up to 1 out of 3 adolescents and adults)
- Fever (1 person out of 10, or less)
- Mild rash, up to a month after vaccination (1 person out of 25). It is possible for these people to infect other members of their household, but this is extremely rare.

Moderate problems

- Seizure (jerking or staring) caused by fever (very rare).

Severe problems

- Pneumonia (very rare)

Other serious problems, including severe brain reactions and low blood count, have been reported after chickenpox vaccination. These happen so rarely experts cannot tell whether they are caused by the vaccine or not. If they are, it is extremely rare.

Note: The first dose of **MMRV** vaccine has been associated with rash and higher rates of fever than MMR and varicella vaccines given separately. Rash has been reported in about 1 person in 20 and fever in about 1 person in 5.

Seizures caused by a fever are also reported more often after MMRV. These usually occur 5–12 days after the first dose.

5**What if there is a serious reaction?****What should I look for?**

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS is only for reporting reactions. They do not give medical advice.

6**The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

7**How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

**Vaccine Information Statement (Interim)
Varicella Vaccine**

3/13/2008

42 U.S.C. § 300aa-26

Office Use Only



Hepatitis A Vaccine

What You Need to Know

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 What is hepatitis A?

Hepatitis A is a serious liver disease caused by the hepatitis A virus (HAV). HAV is found in the stool of people with hepatitis A.

It is usually spread by close personal contact and sometimes by eating food or drinking water containing HAV. A person who has hepatitis A can easily pass the disease to others within the same household.

Hepatitis A can cause:

- “flu-like” illness
- jaundice (yellow skin or eyes, dark urine)
- severe stomach pains and diarrhea (children)

People with hepatitis A often have to be hospitalized (up to about 1 person in 5).

Adults with hepatitis A are often too ill to work for up to a month.

Sometimes, people die as a result of hepatitis A (about 3–6 deaths per 1,000 cases).

Hepatitis A vaccine can prevent hepatitis A.

2 Who should get hepatitis A vaccine and when?

WHO

Some people should be routinely vaccinated with hepatitis A vaccine:

- All children between their first and second birthdays (12 through 23 months of age).
- Anyone 1 year of age and older traveling to or working in countries with high or intermediate prevalence of hepatitis A, such as those located in Central or South America, Mexico, Asia (except Japan), Africa, and eastern Europe. For more information see www.cdc.gov/travel.
- Children and adolescents 2 through 18 years of age who live in states or communities where routine vaccination has been implemented because of high disease incidence.
- Men who have sex with men.
- People who use street drugs.
- People with chronic liver disease.

- People who are treated with clotting factor concentrates.
- People who work with HAV-infected primates or who work with HAV in research laboratories.
- Members of households planning to adopt a child, or care for a newly arriving adopted child, from a country where hepatitis A is common.

Other people might get hepatitis A vaccine in certain situations (ask your doctor for more details):

- Unvaccinated children or adolescents in communities where outbreaks of hepatitis A are occurring.
- Unvaccinated people who have been exposed to hepatitis A virus.
- Anyone 1 year of age or older who wants protection from hepatitis A.

Hepatitis A vaccine is not licensed for children younger than 1 year of age.

WHEN

For children, the first dose should be given at 12 through 23 months of age. Children who are not vaccinated by 2 years of age can be vaccinated at later visits.

For others at risk, the hepatitis A vaccine series may be started whenever a person wishes to be protected or is at risk of infection.

For travelers, it is best to start the vaccine series at least one month before traveling. (Some protection may still result if the vaccine is given on or closer to the travel date.)

Some people who cannot get the vaccine before traveling, or for whom the vaccine might not be effective, can get a shot called immune globulin (IG). IG gives immediate, temporary protection.

Two doses of the vaccine are needed for lasting protection. These doses should be given at least 6 months apart.

Hepatitis A vaccine may be given at the same time as other vaccines.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

3**Some people should not get hepatitis A vaccine or should wait.**

- Anyone who has ever had a severe (life threatening) allergic reaction to a previous dose of hepatitis A vaccine should not get another dose.
- Anyone who has a severe (life threatening) allergy to any vaccine component should not get the vaccine.
- **Tell your doctor if you have any severe allergies**, including a severe allergy to latex. All hepatitis A vaccines contain alum, and some hepatitis A vaccines contain 2-phenoxyethanol.
- Anyone who is moderately or severely ill at the time the shot is scheduled should probably wait until they recover. Ask your doctor. People with a mild illness can usually get the vaccine.
- Tell your doctor if you are pregnant. Because hepatitis A vaccine is inactivated (killed), the risk to a pregnant woman or her unborn baby is believed to be very low. But your doctor can weigh any theoretical risk from the vaccine against the need for protection.

4**What are the risks from hepatitis A vaccine?**

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. The risk of hepatitis A vaccine causing serious harm, or death, is extremely small.

Getting hepatitis A vaccine is much safer than getting the disease.

Mild problems

- soreness where the shot was given (*about 1 out of 2 adults, and up to 1 out of 6 children*)
- headache (*about 1 out of 6 adults and 1 out of 25 children*)
- loss of appetite (*about 1 out of 12 children*)
- tiredness (*about 1 out of 14 adults*)

If these problems occur, they usually last 1 or 2 days.

Severe problems

- serious allergic reaction, within a few minutes to a few hours after the shot (*very rare*).

5**What if there is a serious reaction?****What should I look for?**

- Look for anything that concerns you, such as signs of a severe allergic reaction, very high fever, or behavior changes.

Signs of a severe allergic reaction can include hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, and weakness. These would start a few minutes to a few hours after the vaccination.

What should I do?

- If you think it is a severe allergic reaction or other emergency that can't wait, call 9-1-1 or get the person to the nearest hospital. Otherwise, call your doctor.
- Afterward, the reaction should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your doctor might file this report, or you can do it yourself through the VAERS web site at www.vaers.hhs.gov, or by calling 1-800-822-7967.

VAERS is only for reporting reactions. They do not give medical advice.

6**The National Vaccine Injury Compensation Program**

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines.

Persons who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling 1-800-338-2382 or visiting the VICP website at www.hrsa.gov/vaccinecompensation.

7**How can I learn more?**

- Ask your doctor.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call 1-800-232-4636 (1-800-CDC-INFO) or
 - Visit CDC's website at www.cdc.gov/vaccines

**Vaccine Information Statement (Interim)
Hepatitis A Vaccine**

10/25/2011

42 U.S.C. § 300aa-26

Office Use
Only